

Bermudagrass in Kansas

Bermudagrass has many desirable qualities fitting today's turfgrass needs and environmental concerns. It is the primary turfgrass in the southern United States and is widely used in Kansas. The potential for water savings with bermudagrass has caused a considerable increase in its use.

Bermudagrass has several positive attributes. It is a durable, heat- and drought-resistant warm-season grass that requires less water than most other grasses. This grass forms a medium-green, dense, low-growing turf with a medium-fine texture. Bermudagrass is quick to recover from wear injury; thus, it is well-adapted for heavily used areas. Compared to buffalograss, bermudagrass is equal in heat and drought tolerance and water requirement, but it forms a denser, greener, more wear- and weed-resistant turf. However, it does require more fertilizer and mowing than buffalograss.

Bermudagrass adapts to a wide range of soil types and has a deep, vigorous root system. Creeping stems growing above and below ground form a dense, erosion-resistant sod. This is valuable on hot, south-facing slopes where other grasses often fail. Also, because of its low growth habit, bermudagrass adapts to close mowing in highly maintained areas.

Bermudagrass grows best in hot, sunny areas where cool-season grasses are difficult to maintain. It is not a shade-tolerant grass, but a few hours of shade each day will not result in a noticeable decline in quality. However, as shade increases, bermudagrass will decline in quality and density.

On the negative side, bermudagrass is difficult to keep out of flower beds and gardens because of its aggressive growth habit. Also, because it is a warm-season grass, it greens up later in the spring and turns brown earlier in the fall than cool-season grasses such as tall fescue or Kentucky bluegrass. In addition, bermudagrass

Header photo taken at Children's Mercy Park Stadium in Kansas City, KS

generally has a higher likelihood of winter damage compared to specific cultivars of buffalograss or zoysiagrass. Selection of more cold-hardy cultivars is quite important, and more information is provided in detail below.

Care and Management Summary*

Cultivars	Vegetative types that are winter hardy (establish from sod, plugs, or sprigs): Tahoma 31; Northbridge; Latitude 36 Seed type that is winter hardy (established from seed): Monaco
Planting	Mid-May–July Seed: 1.5–3 lb. / 1,000 sq. ft. Sod: 111 sq. yd. / 1,000 sq. ft. Plugs: 1,000 / 1,000 sq. ft. (3+ yd. of sod required) Sprigs: 5–15 bu. / 1,000 sq. ft.
Mowing	~0.5 inch, cut 3 times a week (golf course tees and fairways) 0.75 – 1 inch, cut 3 times a week (sports fields) 1 – 2 inch for home lawns (cut once or twice weekly)
Watering	Soak soil 6–8 inches deep at first signs of wilting; wait for signs of wilting before watering again. Water only during hot, dry weather.
Fertilizing	May–August, 1 lb. actual nitrogen per 1,000 sq. ft. per application; 2–4 spaced applications depending on level of maintenance desired
Weeds	Control dandelion, chickweed, henbit, and other broadleaf weeds in the fall. Apply pre-emergence herbicide in the spring when redbud trees are reaching full bloom to prevent crabgrass

*See text for more detailed information

Recommended Cultivars

Many bermudagrass cultivars on the market are developed in the South and are not adapted to Kansas. It is important to select a cold-hardy variety that will survive Kansas winters. Several vegetative bermudagrass cultivars have been developed by breeders at Oklahoma State University; details and photos below.

Bermudagrass is sold by sod or seed, depending on the variety. The best bermudagrasses are the vegetative types that must be planted by sod, plugs, or sprigs.

Because of their hybrid breeding, they either produce sterile seed or seed unlike the parent plant.

Most seeded cultivars are produced in southwestern states and do not survive cold Kansas winters. However, the seeded variety Monaco, developed by Johnston Seed Company in Oklahoma, has good cold tolerance and can be grown in Kansas.

Rio bermudagrass is another seeded type provided by Johnston Seed Company, and also has good cold tolerance.



Tahoma 31 (vegetative)



Northbridge (vegetative)



Latitude 36 (vegetative)



Monaco (seeded)

Planting

Because bermudagrass is a warm-season grass, it establishes best after the soil and air temperatures are warm and there is no longer a danger from spring frost. Mid-May through July is the best time for planting. Later plantings may not completely fill in before winter, while earlier plantings are slow to establish and often are taken over by weeds.

The soil should be prepared several weeks before it is time to plant. Soil preparation is the same whether using seed, sod, plugs, or sprigs. The first step is to test the soil to determine if any of the essential nutrients are deficient. Information on how to take soil samples for testing can be obtained at your county Extension office or a private soil testing lab. Prices will vary among labs and also depend upon how extensive a test is requested.

Do not leave low spots where water will stand. After grading, till the soil as deeply as possible (it is often difficult to till deeper than 8 to 12 inches), and incorporate recommended nutrients from the soil test. After tilling and incorporating nutrients, a final finish grading is needed.

Sod is the quickest but most expensive method used to establish bermudagrass. After it is placed onto prepared soil, one has an instant bermudagrass turf. However, frequent watering is needed until the sod roots into the soil. Approximately 111 square yards of sod are required for 1,000 square feet of lawn area.

Sod can be cut into 2 × 2-inch square, or 2-inch diameter round sections called plugs to make it go farther. You may ask your supplier to cut them, because it is a difficult task. Plugs planted 12 inches apart will grow together in one season unless they are planted late.

Weeds likely will grow in the bare soil between the plugs until a dense turf is formed, but can be controlled either by hand removal or herbicides. After weed control, mow the bermudagrass to encourage faster spreading so the plugs will fill in.

Sprigging is a method of planting stems from shredded turf in shallow furrows. This method is more commonly used on sports turf and golf course fairways. Stolonizing is an alternative method where the sprigs are scattered on the soil surface and covered with topdressing.

If using a seeded variety, broadcast the seed on prepared soil with a fertilizer spreader, rake lightly, and water in. Applying a thin layer of weed-free straw over the soil surface will protect the seed and speed up germination.

Seeding rates depend on the kind of seed purchased — hulled, unhulled, or coated.

Suggested Seeding Rates

Form of seed	Seeds/lb.	Seeding rate/ 1,000 sq. ft.
Unhulled	1.6 million	1.5–2.0 lb.
Hulled	2.1 million	1.0–1.5 lb.
Hulled-coated	1.0 million	1.5 –2.0 lb.
Unhulled-coated	less than 1.0 million*	2.5–3.0 lb.

*Variable depending on coating method

With all methods of planting, it will be necessary to keep the soil moist until the grass is established. Keep weeds under control and begin mowing as soon as the grass becomes tall enough to mow.

You can see more details and photos regarding planting options in the extension document “Zoysia-grass in Kansas.”

Maintenance

Mowing, watering, and fertilizing are the fundamental cultural practices forming the foundation of a turf maintenance program. Each cultural practice affects the other. For example, increasing the amount of fertilizer also will increase the amount of mowing and watering required.

You must decide on the quality of bermudagrass you want, as well as the amount of work you are willing to expend on your turf area. Do you want a low-maintenance bermudagrass area or one with more visual appeal? A high bermudagrass area is possible with a reasonable amount of maintenance, but it must be done in a timely and correct fashion. It is important to follow a planned, complete program based on recommended maintenance practices.

Mowing. Select a mowing height based on your maintenance level. Bermudagrass can be mowed anywhere from ½ to 2 inches. When mowed twice a week at 1 inch with a reel mower, bermudagrass forms a dense and even turf. Mowing at 2 inches results in less maintenance, requiring mowing only once every 10 days or so. Also, higher-cut turf does not need to be watered as much and is more resistant to weeds. When mowing, don’t remove more than one-third of the foliage. Cutting too much foliage will put the grass in shock and result in a stemmy turf. The shorter you keep the grass, the more often you will need to mow.

Keep the mower blade sharp so the mower cuts the leaf tips cleanly. This is especially important when mowing a newly planted turf. When mowing higher than 1 inch, use a rotary mower. If you prefer a shorter lawn (1 inch or less) a reel mower will be required. It is not necessary to collect clippings if you mow frequently enough—the clippings will filter into the turf where they quickly decompose and return nutrients to the soil.

Watering. Bermudagrass is drought resistant and does not need as much water as tall fescue. Don’t water on a calendar basis; rather, monitor the turf and the soil. Water when it is hot and dry, watching for signs of wilting. The onset of wilting is accompanied by a change in turf color to a bluish-gray cast.

When watering, soak the turf thoroughly to a depth of 6 to 8 inches. Avoid light, frequent watering (except on newly planted lawns) because it causes shallow rooting. Try to water in the morning when the temperature is cooler and it is less windy.

Fertilizing. The amount of fertilizer used largely determines the level of turf maintenance. If only 2 pounds of actual nitrogen per 1,000 square feet are applied during the season, less watering and mowing will be required. Applying 4 pounds of actual nitrogen fertilizer per 1,000 square feet will produce a dark green color, but more mowing and watering will be required.

Many types of fertilizers are available. The only way to know which formulation is appropriate for your lawn is to have your soil tested. Fertilizer can be applied in either liquid or granular form. Both are effective; the only difference between them is the method of application. There are two types of fertilizer spreaders: rotary spreaders, which throw the fertilizer in a wide pattern, and drop spreaders. The drop method can be more accurate, because it drops the fertilizer between the wheels of the spreader, although it takes longer to cover the lawn. Fertilizer should be applied May through August.

Put only 1 pound of nitrogen fertilizer per 1,000 square feet on bermudagrass at a time; more may cause excessive growth. The grass foliage should be dry at the time of application to avoid fertilizer burn. Be careful when applying fertilizer — ensure the correct amount is uniformly applied. An uneven application will cause streaks in the turf area.

Thatch. Bermudagrass has a tendency to form thatch.

Thatch is a layer of decomposing organic matter that looks somewhat like peat moss. It is located between the soil surface and the grass foliage and is made up of surface roots, stems, and runners. Thatch buildup occurs when organic matter is produced faster than soil microorganisms can decompose it.

Thatch can be beneficial or detrimental to the lawn. When thatch is excessive, movement of water, nutrients, and pesticide into the soil can be restricted. Roots form in the thatch instead of penetrating deep into the soil, making the turf less drought resistant. Thatch also provides a habitat for disease organisms and insects.

To determine the amount of thatch in bermudagrass, cut a small wedge of turf down into the soil with a knife. If the layer is greater than ½ inch, the thatch should be controlled. The best times for thatch control are June or July when the bermudagrass is actively growing.

There are two types of power equipment for thatch control — core aerifiers and power rakes. Equipment can be rented, or you can hire a professional service. A core aerator is a machine that pulls cores out of the soil.

These cores are usually about ¾ inch in diameter, 2 to 3 inches deep, and about 3 inches apart. The plugs are beneficial and should be left to decompose. Decomposition of the plugs could take several weeks, depending on the type of soil. Alternatively, the thatch can be removed by power raking. However, this method is more destructive to the turf, and produces debris that will have to be removed.

Newly established bermudagrass will not have thatch, as it takes several years for it to build up. But once it is established, there are some measures one can take to prevent thatch. Avoid applying excessive amounts of nitrogen fertilizer. Turf that is growing faster produces more thatch. Water as deeply and infrequently as possible, without stressing the grass. This helps produce a deeper root system; shallow-rooted lawns develop more thatch.

Weeds. Bermudagrass is fairly resistant to weeds; however there are occasional problems if the turf is thin. Weed invasion of bermudagrass occurs primarily in early spring and late fall when the grass is dormant and unable to compete with the weeds. The most common weed problems are dandelion, chickweed, and henbit. It is best to control these weeds in the fall when they are young. Use herbicide labeled for the particular weed being sprayed. By spring, weeds are established and more difficult to control. Additionally, chickweed and henbit are winter annuals that will die in the summer when bermudagrass is vigorously growing. Therefore, spring applications of herbicides to those weeds are wasteful.

Bermudagrass Maintenance Calendar

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Planting						■	■	■				
Mowing				■	■	■	■	■	■			
Watering					■	■	■	■	■			
Fertilizing					■	■	■	■				
Dethatching						■	■					
Aerifying					■	■	■	■				
Crabgrass control			■	■	■							
Dandelion control									■	■		

Spring weed control largely consists of preventing crabgrass by using a preemergence herbicide. This should be applied at about the time redbud trees are in full bloom, or are beginning to leaf out.

Unknown weeds should be taken to your county Extension office for identification so you can purchase the correct herbicide and apply it at the right time.

Insects. Bermudagrass is not particularly susceptible to insect problems. As with all turfgrasses, however, grubs can be a problem. If unsure of an insect, have it identified to ensure proper control. Pesticides should not be routinely applied for insect control; use them only when insects are actually damaging the turf. K-State Research and Extension has several publications available on turfgrass insect control.

Disease. The main disease concern with bermudagrass is a fungal disease called spring dead spot. This disease causes round patches of dead grass 6 inches to 1 foot in diameter in the spring. The best prevention is to use a variety that has some resistance to this fungus. The more cold-hardy cultivars seem to be the most resistant, although they can be affected. To prevent other diseases, maintain a healthy turf by following proper watering, fertilizing, and mowing techniques.

Remember, when using pesticides follow all safety and application instructions on the label.

Jack Fry, Professor and Commercial Turf Extension Specialist
Department of Horticulture and Natural Resources, Kansas State University

Steve Keeley, PhD, Department Head
Turfgrass Science | Horticultural Pest Management, Kansas State University



Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

Publications from Kansas State University are available at bookstore.ksre.ksu.edu.

Date shown is that of publication or last revision. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. In each case, credit Jack Fry and Steve Keeley, *Bermudagrass in Kansas*, Kansas State University, August 2025.

**Kansas State University Agricultural Experiment Station
and Cooperative Extension Service**

K-State Research and Extension is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of K-State Research and Extension, Kansas State University, County Extension Councils, Extension Districts.

MF1112 rev. August 2025